



A Conjecture of S. Chowla Via the Generalized Riemann Hypothesis

R. A. Mollin; H. C. Williams

Proceedings of the American Mathematical Society, Vol. 102, No. 4 (Apr., 1988),
794-796.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28198804%29102%3A4%3C794%3AACOSCV%3E2.0.CO%3B2-I>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

A CONJECTURE OF S. CHOWLA VIA THE GENERALIZED RIEMANN HYPOTHESIS

R. A. MOLLIN AND H. C. WILLIAMS

(Communicated by Larry J. Goldstein)

ABSTRACT. S. Chowla conjectured that if $p = m^2 + 1$ is prime and $m > 26$, then h_K , the class number of $K = \mathbb{Q}(\sqrt{p})$, is greater than 1. We prove this conjecture under the assumption of the Riemann hypothesis for ζ_K , the zeta function of K , i.e. the generalized Riemann hypothesis (GRH).

It is the purpose of this note to prove the following result.

THEOREM. *Let $K = \mathbb{Q}(\sqrt{p})$, where $p = m^2 + 1$ is prime and $m > 26$. If the Riemann hypothesis holds for ζ_K then $h_K > 1$.*

Without the GRH hypothesis, this is known as the Chowla conjecture given in [1]. We note that it is an easy consequence of the celebrated Brauer-Siegel theorem that there are only finitely many such p for which $h_K = 1$. In [3] Mollin reduced the problem to the case where $m = 2r$ and $r > 13$ is prime.

In what follows we make use of an idea of Cornell and Washington [2] to show how to use the GRH to get an effective bound ($p > 10^{23}$) for which the Chowla conjecture holds. The remaining finite cases are then handled by a simple sieve process. Throughout the remainder of the paper p will denote a prime of the form $4r^2 + 1$ where r is an odd prime, and K will denote $\mathbb{Q}(\sqrt{p})$.

The following result contains facts which are either well known or trivial. Therefore we state it without proof.

LEMMA. *Let y be a real number.*

- (1) *If $y > 1$, then $\sum 1/q < \log y$, where the sum ranges over all primes $q \leq y$.*
- (2) *If $y > 0$, then $\sum (1/q^2) < 1/y$, where the sum ranges over all primes $q > y$,*
- (3) *If $|y| \leq 1/2$, then $|\log(1 - y) + y| < y^2$.*

Now we are in a position to prove the Theorem.

Let $\varepsilon(q) = (p/q)$ denote the Kronecker symbol where q is a prime. Set

$$T_1(y) = \sum_{q \leq y} [q/(q - \varepsilon(q))] \quad \text{and} \quad T_2(y) = \prod_{q > y} [q/(q - \varepsilon(q))].$$

Received by the editors August 13, 1986 and, in revised form, December 8, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 12A50, 12A25; Secondary 10B05.

Key words and phrases. Class number 1, real quadratic field, Riemann hypothesis.

The first author's research is supported by N.S.E.R.C. Canada, Grant #A8484.

The second author's research is supported by N.S.E.R.C. Canada, Grant #A7649.

Therefore $L(1, \chi) = T_1(y)T_2(y)$, the Euler product for the Dirichlet L -function. Clearly we have that $T_1(y) \geq \prod_{q \leq y} (q/(q+1))$. Therefore

$$(i) \quad \log T_1(y) \geq - \sum_{q \leq y} \log(q+1/q) > - \sum_{q \leq y} 1/q > -\log y,$$

where the last inequality is from Lemma (1). Furthermore, we have from Lemma, (2)–(3), that

$$(ii) \quad \log T_2(y) = - \sum_{q > y} \log(1 - (\varepsilon(q)/q)) > \sum_{q > y} \varepsilon(q)/q - 1/y.$$

By similar reasoning to that used by Cornell and Washington [2, p. 265] (where the GRH is assumed) we get

$$(iii) \quad \sum_{q > y} \varepsilon(q)/q \geq -B(y)[(4 + 3 \log y)/\sqrt{y}],$$

where $B(y) = (\log p)((1/\pi \log y) + (5.3/(\log y)^2)) + 4/\log y + 1/\pi$. Now, from (ii) and (iii), we get

$$(iv) \quad \log T_2(y) > -B(y)[(4 + 3 \log y)/\sqrt{p}] - 1/y.$$

Hence from (i) and (iv) we have

$$L(1, \chi) = T_1(y)T_2(y) > (1/y) \exp[-B(y)[(4 + 3 \log y)/\sqrt{p}] - 1/y].$$

Set $y = (\log p)^2$; whence $\log y = 2 \log \log p$. Thus

$$\begin{aligned} & -B(y)[(4 + 3 \log y)/\sqrt{y}] - 1/y \\ &= - \{2/(\pi \log \log p) + 15/(\log \log p)^2 + 8/(\log \log p) \log p + 4/(\pi \log p) \\ & \quad + 3/\pi + 135/(2 \log \log p) + 12/\log p + 6 \log \log p/(\pi \log p) + 1/(\log p)^2\} \\ &= d(p), \end{aligned}$$

say. Hence $L(1, \chi) > e^{d(p)}/(\log p)^2$. However, $2h_K R = \sqrt{p}L(1, \chi)$, where $R = \log(2r + \sqrt{p}) < \log 2\sqrt{p}$ is the regulator of K ; whence

$$h_K > \frac{\sqrt{p}e^{d(p)}}{2(\log p)^2(\log 2\sqrt{p})} = F(p),$$

say. Moreover $-d(p)$ is a decreasing function of p so then $e^{d(p)}$ is an increasing function of p , forcing $F(p)$ to be an increasing function of p . Thus, if $p > 10^{13}$ then $d(p) < 4.68$, $e^{d(p)} > 0.009279493$ and $\sqrt{p}/(2(\log p)^2(\log 2\sqrt{p})) \geq 112.6838154$. Hence $F(p) > 1$ for $p > 10^{13}$.

Now, to deal with the primes $p < 10^{13}$ where $p = 4r^2 + 1$ we note that $r < 10^{6.5}/2 \approx 1.6 \times 10^6$. Hereafter, we use the fact proved by Mollin in [3] that $h_K = 1$ is tantamount to all primes less than r being inert in K . Select some positive integer k and the first k primes $\{q_i\}_{i=1}^k$ with $q_1 = 5$. For each of these q_i find (by trial) and tabulate those $S_{i,j}$ such that $0 \leq S_{i,j} \leq q_i - 1$ and $((4S_{i,j}^2 + 1)/q_i) \not\equiv -1$ where $(/)$ is the Legendre symbol. There are approximately $q_i/2$ of these. If any $r \equiv S_{i,j} \pmod{q_i}$ and $r > q_i$ then delete this value of r since either: $4r^2 + 1 \equiv 0 \pmod{q_i}$, which means that $4r^2 + 1$ is not prime, or $((4r^2 + 1)/q_i) = 1$ and $q_i < r$, which means that $h_K > 1$. Since half of the r 's are eliminated for each q_i , then a

value of k such that $2^k > 1.6 \times 10^6$ should suffice to complete the task. We used $k = 100$ and a Fortran program to sieve out as many values of $r < 1.6 \times 10^6$ as possible. In a matter of a few minutes we found that if $r > 13$ and $p = 4r^2 + 1$ is a prime ($p < 10^{13}$) there exists some $q_i < r$ such that q_i is quadratic residue of p . This proves the Theorem. Q.E.D.

REFERENCES

1. S. Chowla and J. Friedlander, *Class numbers and quadratic residues*, Glasgow Math. J. **17** (1976), 47–52.
2. G. Cornell and L. C. Washington, *Class numbers of cyclotomic fields*, J. Number Theory **21** (1985), 260–274.
3. R. Mollin, *Necessary and sufficient conditions for the class number of a real quadratic field to be one and a conjecture of S. Chowla*, Proc. Amer. Math. Soc. **102** (1988), 17–21.

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF CALGARY,
2500 UNIVERSITY DRIVE N.W., CALGARY, ALBERTA, CANADA T2N 1N4

COMPUTER SCIENCE DEPARTMENT, THE UNIVERSITY OF MANITOBA, WINNIPEG,
MANITOBA, CANADA R3T 2N2